

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 April 2002 (04.04.2002)

PCT

(10) International Publication Number
WO 02/26705 A2

- (51) International Patent Classification⁷: C07D 207/00 (72) Inventors: BOAZ, Neil, Warren; 1342 Watauga Street, Kingsport, TN 37660 (US). DEBENHAM, Sheryl, Davis; 2 Blue Ridge Circle, Scotch Plains, NJ 7076 (US).
- (21) International Application Number: PCT/US01/30665
- (22) International Filing Date: 28 September 2001 (28.09.2001) (74) Agent: BLAKE, Michael, J.; P.O. Box 511, Kingsport, TX 37662-5075 (US).
- (25) Filing Language: English (81) Designated State (national): JP.
- (26) Publication Language: English (84) Designated States (regional): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).
- (30) Priority Data:
00/236,564 29 September 2000 (29.09.2000) US
60/264,411 26 January 2001 (26.01.2001) US
09/957,182 20 September 2001 (20.09.2001) US
- Published:
— without international search report and to be republished upon receipt of that report
- (71) Applicant: EASTMAN CHEMICAL COMPANY [US/US]; 100 North Eastman Road, Kingsport, TN 37660 (US). For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/26705 A2

(54) Title: HIGHLY ENANTIOMERICALLY PURE LACTAM-SUBSTITUTED PROPANOIC ACID DERIVATIVES AND METHODS OF MAKING AND USING SAME

(57) Abstract: The present invention relates to highly enantiomerically pure lactam-substituted propanoic acid derivatives and methods of making and using therefor. The invention involves a multi-step synthesis to produce the lactam compounds. In one step of the reaction sequence, asymmetric hydrogenation of a lactam-enamide was performed to produce an intermediate that can ultimately be converted to a series of pharmaceutical compounds. The invention also contemplates the *in situ* synthesis of an intermediate of the multi-step synthesis, which provides economic advantages to the overall synthesis of the lactam compounds.

**HIGHLY ENANTIOMERICALLY PURE
LACTAM-SUBSTITUTED PROPANOIC ACID DERIVATIVES AND
METHODS OF MAKING AND USING SAME**

5 CROSS REFERENCE TO RELATED APPLICATIONS

This application claims priority to United States Provisional Application
Nos. 60/236,564, filed September 29, 2000, and 60/264,411, filed January 26,
2001, both entitled "Phosphino-Aminophosphines," which applications are
10 hereby incorporated by this reference in their entireties.

BACKGROUND OF THE INVENTION

Asymmetric catalysis is the most efficient method for the generation of
15 products with high enantiomeric purity, as the asymmetry of the catalyst is
multiplied many times over in the generation of the chiral product. These chiral
products have found numerous applications as building blocks for single
enantiomer pharmaceuticals as well as in some agrochemicals. The asymmetric
catalysts employed can be enzymatic or synthetic in nature. The latter types of
20 catalyst have much greater promise than the former due to much greater latitude
of applicable reaction types. Synthetic asymmetric catalysts are usually
composed of a metal reaction center surrounded by an organic ligand. The
ligand usually is generated in high enantiomeric purity, and is the agent inducing
the asymmetry. A prototypical reaction using these types of catalyst is the
25 asymmetric hydrogenation of enamides to afford amino-acid derivatives
(Ohkuma, T.; Kitamura, M.; Noyori, R. In *Catalytic Asymmetric Synthesis*, 2nd
ed.; Ojima, I., Ed.; Wiley-VCH: New York, 2000; pp. 1-17).

Although the preparation of enamides through Horner-Emmons Wittig
30 chemistry is known, the preparation and use of substrates such as lactam-
substituted 2-propenoic acid derivatives which possess a fully substituted
nitrogen on the enamide are not known and the viability of the standard
preparative sequence for the enamide is unclear. In general, the majority of

enamides that have undergone asymmetric hydrogenation possess a hydrogen substituent on the nitrogen of the enamide. Thus the efficacy of asymmetric catalysts for the hydrogenation of lactam-substituted 2-propenoic acid derivatives is also unclear.

5

U.S. Patent No. 4,696,943 discloses the synthesis of single enantiomer lactam-substituted propanoic acid derivatives useful as pharmaceutical agents for various conditions. However, these compounds were prepared by a cyclization reaction and not by the asymmetric hydrogenation of an enamide.

10

In light of the above, it would be desirable to produce single enantiomer lactam-substituted propanoic acid derivatives useful as pharmaceutical compounds.

15 SUMMARY OF THE INVENTION

The present invention relates to highly enantiomerically pure lactam-substituted propanoic acid derivatives and methods of making and using therefor. The invention involves a multi-step synthesis to produce the lactam
20 compounds. In one step of the reaction sequence, asymmetric hydrogenation of a lactam-enamide was performed to produce an intermediate that can ultimately be converted to a series of pharmaceutical compounds. The invention also contemplates the *in situ* synthesis of an intermediate of the multi-step synthesis, which provides economic advantages to the overall synthesis of the lactam
25 compounds.

Additional advantages of the invention will be set forth in part in the description that follows, and in part will be apparent from the description or may be learned by practice of the invention. The advantages of the invention
30 will be realized and attained by means of the elements and combinations particularly pointed out in the appended claims. It is to be understood that both the foregoing general description and the following detailed description are

exemplary and explanatory only and are not restrictive of the invention, as claimed.

DETAILED DESCRIPTION OF THE INVENTION

5

The present invention may be understood more readily by reference to the following detailed description of aspects of the invention and the Examples included therein.

10 Before the present compositions of matter and methods are disclosed and described, it is to be understood that this invention is not limited to specific synthetic methods or to particular formulations, and, as such, may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular aspects only and is not intended to be limiting.

15

The singular forms a, an, and the include plural referents unless the context clearly dictates otherwise.

Optional or optionally means that the subsequently described event or
20 circumstances may or may not occur, and that the description included instances where said event or circumstance occurs and instances where it does not.

The term "alkyl group" may include straight- or branched-chain, aliphatic hydrocarbon radicals containing up to 20 carbon atoms and may be
25 substituted, for example, with one to three groups selected from C₁-C₆-alkoxy, cyano, C₂-C₆-alkoxycarbonyl, C₂-C₆ alkanoyloxy, hydroxy, aryl and halogen. The terms "C₁-C₆-alkoxy," "C₂-C₆-alkoxycarbonyl," and "C₂-C₆-alkanoyloxy" are used to denote radicals corresponding to the structures -OR, -CO₂R, and -OCOR, respectively, wherein R is C₁-C₆-alkyl or substituted C₁-C₆-alkyl.

30

The term "cycloalkyl" is used to denote a saturated, carbocyclic hydrocarbon. The term "substituted cycloalkyl" is a cycloalkyl group

substituted with one or more of the groups described above.

The term "aryl group" may include phenyl, naphthyl, or anthracenyl and phenyl, naphthyl, or anthracenyl substituted with one to three substituents
5 selected from C₁-C₆-alkyl, substituted C₁-C₆-alkyl, C₆-C₁₀ aryl, substituted C₆-C₁₀ aryl, C₁-C₆-alkoxy, halogen, carboxy, cyano, C₁-C₆-alkanoyloxy, C₁-C₆-alkylthio, C₁-C₆-alkylsulfonyl, trifluoromethyl, hydroxy, C₂-C₆-alkoxycarbonyl, C₂-C₆-alkanoylamino and -OR', SR', -SO₂R', -NHSO₂R' and -NHCO₂R',
10 wherein R' is phenyl, naphthyl, or phenyl or naphthyl substituted with one to three groups selected from C₁-C₆-alkyl, C₆-C₁₀ aryl, C₁-C₆-alkoxy and halogen.


The term "heteroaryl group" includes a 5- or 6- membered aromatic ring containing one to three heteroatoms selected from oxygen, sulfur and nitrogen. Examples of such heteroaryl groups are thienyl, furyl, pyrrolyl, imidazolyl,
15 pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, triazolyl, thiadiazolyl, oxadiazolyl, tetrazolyl, pyridyl, pyrimidyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, indolyl and the like. The heteroaryl group may be substituted, for example, with up to three groups such as C₁-C₆-alkyl, C₁-C₆-alkoxy, substituted C₁-C₆-alkyl, halogen, C₁-C₆-alkylthio, aryl, arylthio, aryloxy, C₂-C₆-
20 alkoxycarbonyl and C₂-C₆-alkanoylamino. The heteroaryl group also may be substituted with a fused ring system, e.g., a benzo or naphtho residue, which may be unsubstituted or substituted, for example, with up to three of the groups set forth in the preceding sentence.

25 Reference will now be made in detail to the present aspects of the invention. Wherever possible, the same reference numbers and letters are used throughout the various formulas in the invention to refer to the same or like parts.


30 The present invention relates to the synthesis of enantiomerically pure lactam-substituted propanoic acid derivatives and methods of making and using

therefor. A reaction scheme that depicts a general sequence of reaction steps to produce the compounds of the invention is shown in Scheme 1.

SCHEME 1




The first step depicted in Scheme 1 involves the reaction (*i.e.*, condensation) between a compound having the formula I



with glyoxylic acid, wherein R¹ is hydrogen, substituted or unsubstituted, branched or straight chain C₁ to C₂₀ alkyl; substituted or unsubstituted C₃ to C₈ cycloalkyl; substituted or unsubstituted C₆ to C₂₀ aryl; or substituted or unsubstituted C₄ to C₂₀ heteroaryl, and

n is from 0 to 5,


to produce a compound having the formula II.



The condensation reaction between lactam I and glyoxylic acid is generally conducted in a solvent. Examples of useful solvents include, but are not limited to, aliphatic hydrocarbons such as hexane, heptane, octane and the like, aromatic hydrocarbons such as toluene, xylenes, and the like, cyclic or acyclic ethers such as diethyl ether, tert-butyl methyl ether, diisopropyl ether, tetrahydrofuran and the like, or polar aprotic solvents such as dimethylformamide, dimethyl sulfoxide, N-methylpyrrolidone and the like. The amount of glyoxylic acid relative to the amount of compound I can vary. In one aspect, the glyoxylic acid is present in the amount from 0.8 to 2 equivalents per 1.0 equivalent of the compound having the formula I. The condensation reaction is generally run

between ambient temperature and the boiling point of the lowest boiling component of the mixture.

The second step depicted in Scheme 1 involves converting compound II
5 to a compound having the formula III




wherein R^1 , R^3 , and R^4 are, independently, a substituted or unsubstituted,
10 branched or straight chain C_1 to C_{20} alkyl group; a substituted or unsubstituted C_3 to C_8 cycloalkyl group; a substituted or unsubstituted C_6 to C_{20} aryl group; or a substituted or unsubstituted C_4 to C_{20} heteroaryl group, wherein R^1 can also be hydrogen, and n is from 0 to 5.

15 The second step generally involves reacting a compound having the formula II with an alcohol comprising an alkyl alcohol, an aryl alcohol, or a heteroaryl alcohol, wherein the alkyl alcohol is substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl or substituted or unsubstituted C_3 to C_8 cycloalkyl; the aryl alcohol is substituted or unsubstituted C_6 to C_{20} aryl; and
20 the heteroaryl alcohol is substituted or unsubstituted C_4 to C_{20} heteroaryl, wherein the heteroatom is oxygen, nitrogen, or sulfur. Although R^3 and R^4 need not be the same group, they can be derived from the same alcohol. In one aspect, the alcohol is a C_1 to C_5 alcohol, preferably methanol or ethanol. The amount of alcohol that is used can vary. In one aspect, the alcohol is present in
25 the amount from 2.0 to 5.0 equivalents per 1.0 equivalent of the compound having the formula II.

In another aspect, the second step is generally conducted under dehydrating conditions using acid catalysis. For example, the use of a dehydrating agent such as a trialkyl orthoformate or the physical removal of water from the reaction mixture via an azeotropic distillation are useful in the present invention. When a dehydrating agent is used, the amount of dehydrating agent used relative to the amount of compound II is generally between 2 and 5 molar equivalents. In general, the alcohol can be used as the reaction solvent for the preparation of compound III; however, co-solvents can be used. Examples of co-solvents useful in the present invention include, but are not limited to, aliphatic hydrocarbons such as hexane, heptane, octane and the like, aromatic hydrocarbons such as toluene, xylenes, and the like. A preferable co-solvent is toluene or xylene. The second step is generally performed at a temperature between ambient temperature and the boiling point of the lowest boiling component of the reaction mixture. In one aspect, the reaction is performed between 20 °C and 120 °C.

The compounds produced in steps 1 and 2 can be represented by the general formula XI




wherein R^1 , R^3 , and R^4 are, independently, hydrogen, a substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl group; a substituted or unsubstituted C_3 to C_8 cycloalkyl group; a substituted or unsubstituted C_6 to C_{20} aryl group; or substituted or unsubstituted C_4 to C_{20} heteroaryl group, and n is from 0 to 5. In one aspect, n is 2. In another aspect, R^1 , R^3 , and R^4 are hydrogen and n is 2. In a further aspect, R^1 is hydrogen, R^3 and R^4 are methyl,

and n is 2.

The third step in Scheme 1 involves converting compound III to the halogenated lactam compound having the formula IV


5



wherein R¹ and R³ are, independently, substituted or unsubstituted, branched or straight chain C₁ to C₂₀ alkyl; substituted or unsubstituted C₃ to C₈ cycloalkyl; substituted or unsubstituted C₆ to C₂₀ aryl; or substituted or unsubstituted C₄ to C₂₀ heteroaryl, wherein R¹ can also be hydrogen, X is fluoride, chloride, bromide, or iodide, and n is from 0 to 5.

The third step generally involves reacting compound III with a phosphorous trihalide having the formula PX₃, wherein X is fluoro, chloro, bromo, or iodo. In one aspect of the invention, the phosphorous trihalide is phosphorous trichloride or phosphorous tribromide. The amount of the phosphorus trihalide is generally between 0.8 and 2.0 molar equivalents based on compound III. Typically, the reaction is performed in an inert solvent including, but not limited to, aliphatic hydrocarbons such as hexane, heptane, octane and the like, aromatic hydrocarbons such as toluene, xylenes, and the like, and halogenated hydrocarbons such as dichloromethane, dichloroethane, tetrachloroethylene, chlorobenzene, and the like. The third step is generally performed at a temperature between ambient temperature and the boiling point of the lowest boiling component of the reaction mixture for a time necessary to consume the majority of compound III. In one aspect, the reaction solvent is toluene or xylene and the reaction temperature is between 40 °C and 80 °C.

The fourth step in Scheme 1 involves converting compound IV to a compound having the formula V



5

wherein R^1 and R^3 are, independently, substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl; substituted or unsubstituted C_3 to C_8 cycloalkyl; substituted or unsubstituted C_6 to C_{20} aryl; or substituted or unsubstituted C_4 to C_{20} heteroaryl, wherein R^1 can also be hydrogen,

10

R^6 is substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl or substituted or unsubstituted C_3 to C_8 cycloalkyl, and

n is from 0 to 5. In one aspect, n is 2 and R^1 is hydrogen. In another aspect of the invention, R^3 is methyl or ethyl. In a further aspect, R^6 is methyl or ethyl.

15

The fourth step is an Arbuzov reaction comprising reacting a compound having the formula IV with a phosphite having the formula $P(OR^6)_3$, wherein R^6 is substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl or substituted or unsubstituted C_3 to C_8 cycloalkyl. In one aspect, the phosphite is trimethyl phosphite or triethyl phosphite. In another aspect of the invention, compound IV is the chloro or bromo compound ($X = Cl$ or Br). The amount of the phosphite is generally between 0.8 and 1.2 molar equivalents based on compound IV. The reaction is optionally conducted in the presence of a solvent including, but not limited to, aliphatic hydrocarbons such as hexane, heptane, octane and the like, aromatic hydrocarbons such as toluene, xylenes, and the


20

25

like, cyclic or acyclic ethers such as tert-butyl methyl ether, diisopropyl ether, tetrahydrofuran and the like, halogenated hydrocarbons such as dichloromethane, dichloroethane, tetrachloroethylene, chlorobenzene and the like, or polar aprotic solvents such as dimethylformamide, dimethyl sulfoxide, N-methylpyrrolidone and the like. The reaction is generally performed at a temperature between ambient temperature and the boiling point of the lowest boiling component of the reaction mixture for a time necessary to consume the majority of compound IV. In one aspect, the solvent is toluene or xylene and the reaction is performed between 40 °C and 100 °C.

10

The fifth step in Scheme 1 involves converting compound V to a compound having the formula VI



15

wherein R^1 , R^2 , and R^3 are, independently, hydrogen, substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl; substituted or unsubstituted C_3 to C_8 cycloalkyl; substituted or unsubstituted C_6 to C_{20} aryl; or substituted or unsubstituted C_4 to C_{20} heteroaryl, and n is from 0 to 5. In one aspect, n is 2 and R^1 is hydrogen. In another aspect, R^2 and R^3 are methyl. In a further aspect, R^2 is methyl and R^3 is ethyl.

The fifth step of the sequence is a Horner-Emmons Wittig reaction between phosphonate compound V and an aldehyde having the formula $HC(O)R^2$ to afford enamide VI. In one aspect of the invention, the aldehyde is acetaldehyde. The reaction generally involves the use of a base. For example, the base can be a moderately strong non-hydroxide base with a pK_a of 13 or


25

above. Examples of non-hydroxide bases include, but are not limited to, amidine bases such as 1,5-diazabicyclo[4.3.0]non-5-ene (DBN), 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), or guanidine bases such as tetramethylguanidine (TMG). The amount of base is usually between 1.0 and 2.0 molar equivalents based on compound V, and the amount of aldehyde is generally between 0.8 and 1.5 molar equivalents based compound V.

The reaction to produce compound VI is generally conducted in the presence of a solvent. Solvents useful in the reaction include, but are not limited to, aliphatic hydrocarbons such as hexane, heptane, octane and the like, aromatic hydrocarbons such as toluene, xylenes, and the like, cyclic or acyclic ethers such as tert-butyl methyl ether, diisopropyl ether, tetrahydrofuran and the like, halogenated hydrocarbons such as dichloromethane, dichloroethane, tetrachloroethylene, chlorobenzene and the like, or polar aprotic solvents such as dimethylformamide, dimethyl sulfoxide, N-methylpyrrolidone and the like. The reaction is generally performed at a temperature between - 80 °C and the boiling point of the lowest boiling component of the reaction mixture for a time necessary to largely consume compound V. In one aspect of the invention, the reaction is performed in toluene or xylene at a temperature of 0 °C to 50 °C.

The invention also contemplates producing compound VI from compound III *in situ*. The term "*in situ*" is defined herein as performing two or more reaction sequences without isolating any of the intermediates that are produced during the reaction sequence. One aspect of the invention involves a method for producing the compound VI *in situ*, comprising

- (a) reacting a compound having the formula III



wherein R^3 and R^4 are a substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl group; a substituted or unsubstituted C_3 to C_8 cycloalkyl group; a substituted or unsubstituted C_6 to C_{20} aryl group; or substituted or unsubstituted C_4 to C_{20} heteroaryl group, wherein R^1 can also be hydrogen,

with PX_3 , wherein X is fluoride, chloride, bromide, or iodide, to produce a halogenated lactam;

(b) reacting the halogenated lactam produced in step (a) with a phosphite having the formula $P(OR^6)_3$, wherein R^6 is substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl or substituted or unsubstituted C_3 to C_8 cycloalkyl, to produce a phosphonated lactam; and


(c) reacting the phosphonated lactam produced in step (b) with an aldehyde having the formula $HC(O)R^2$, wherein R^2 is hydrogen, substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl; substituted or unsubstituted C_3 to C_8 cycloalkyl; substituted or unsubstituted C_6 to C_{20} aryl; or substituted or unsubstituted C_4 to C_{20} heteroaryl,

in the presence of a base,

wherein steps (a), (b), and (c) are performed *in situ*. In this aspect of the invention, compounds IV and V are not isolated. The *in situ* preparation of compound VI would not have been expected due to the incompatibility of several of the reagents used in the *in situ* process. In addition, there is an

economic advantage to combining multiple reaction steps without having to isolate each of compound that is produced.

The sixth step in Scheme 1 involves converting compound VI to a
5 compound having the formula VII



wherein R¹, R², and R³ are, independently, hydrogen, a substituted or
10 unsubstituted, branched or straight chain C₁ to C₂₀ alkyl group; a substituted or
unsubstituted C₃ to C₈ cycloalkyl group; a substituted or unsubstituted C₆ to C₂₀
aryl group; or substituted or unsubstituted C₄ to C₂₀ heteroaryl group,
n is from 0 to 5, and
15 the stereochemistry at carbon a is substantially R or S.

The term “substantially R or S” refers to the enantiomeric purity of the
aforementioned compound, which is measured by the enantiomeric excess of this
20 material. The enantiomeric excess (ee) is defined as the percent of one
enantiomer of the mixture minus the percent of the other enantiomer.
“Substantially R” indicates an ee of 90% or greater with the R enantiomer the
major enantiomer, whilst “substantially S” indicates an ee of 90% or greater
with the S enantiomer the major enantiomer. In one aspect of the invention, the
25 stereochemistry at carbon a is substantially S. In another aspect of the
invention, the stereochemistry at carbon a is substantially R.


In other aspects of the invention, the heteroatom of the heteroaryl group
in compound VII is oxygen, sulfur, or nitrogen, and the substituent on the

substituted alkyl, aryl, or heteroaryl group comprises alkyl, aryl, hydroxy, alkoxy, fluoro, chloro, bromo, iodo, nitro, cyano, or an ester; R² and R³ are independently selected from methyl or ethyl; R¹ is hydrogen, R² is methyl, R₃ is methyl or ethyl; and n is 2.

5

The conversion of compound VI to compound VII comprises hydrogenating compound VI with hydrogen in the presence of a catalyst comprised of a chiral ligand/metal complex to asymmetrically hydrogenate the carbon-carbon double bond of compound VI. The term “hydrogenate” generally refers to reacting a carbon-carbon double or triple bond with hydrogen to reduce the degree of unsaturation. For example, the carbon-carbon double bond of compound VI is hydrogenated to produce a carbon-carbon single bond. Here, the degree of unsaturation has been reduced by one. Referring to Scheme 2, hydrogen can be added to side b (front side) or c (back side) of the carbon-carbon double bond of compound VII. The stereochemistry at carbon a of compound VII will be determined by which side of the carbon-carbon double bond hydrogen approaches. The term “asymmetrically hydrogenating” refers to the addition of hydrogen to a particular side or face (b or c) of the carbon-carbon double bond of compound VII in preference to the other side. The degree of asymmetric hydrogenation is described by the enantiomeric excess of the asymmetric hydrogenation product.

SCHEME 2



25


The asymmetric hydrogenation of compound VI involves the use of a

chiral ligand/metal complex. The chiral ligand/metal complex is composed of a chiral ligand and a metal, where the metal is either chemically bonded to the chiral ligand or the metal is coordinated to the chiral ligand. Any chiral ligand/metal complex known in the art can be used to asymmetrically hydrogenate compound VI. For example, the chiral ligand/metal complexes disclosed in Ohkuma *et al.* in *Catalytic Asymmetric Synthesis*, 2nd Ed, Wiley-VCH, 2000, pages 1-17, which is incorporated by reference in its entirety, are useful in the present invention.

10 In one aspect of the invention, the chiral ligand of the chiral ligand/metal complex comprises the substantially pure enantiomer or diastereomer of 2,3-*O*-isopropylidene-2,3-dihydroxy-1,4-*bis*(diphenylphosphino)butane; 2,2'-*bis*(diphenylphosphino)-1,1'-binaphthyl; 1,2-*bis*-2,5-dialkylphospholano(benzene); 1,2-*bis*-2,5-dialkylphospholano(ethane); 2,3-*bis*-
15 (diphenylphosphino)butane; or 2-diphenylphosphinomethyl-4-diphenylphosphino-1-*t*-butoxycarbonylpyrrolidine.

In another aspect of the invention, the chiral ligand of the chiral ligand/metal complex comprises a substantially enantiomerically pure bis-
20 phosphine compound comprising one phosphine residue having three phosphorus-carbon bonds and the other having two phosphorus-carbon bonds and one phosphorus-nitrogen bond. In one aspect of the invention, the chiral ligand of the chiral ligand/metal complex comprises a phosphine or a bis-phosphine compound and the metal of the chiral ligand/metal complex
25 comprises rhodium, ruthenium, or iridium.

Examples of substantially enantiomerically pure bis-phosphine compounds, *e.g.*, an enantiomeric excess of 90% or greater, include phosphinometallocenyl-aminophosphines having the general formulas IX and X
30 (the enantiomer of IX):



where R^7 , R^8 , R^9 , R^{10} , R^{11} , and R^{12} are, independently, hydrogen, substituted or unsubstituted branched or straight chain C_1 to C_{20} alkyl, substituted or unsubstituted C_3 to C_8 cycloalkyl, substituted or unsubstituted C_6 to C_{20} aryl, and substituted or unsubstituted C_4 to C_{20} heteroaryl, where the heteroatoms are chosen from sulfur, nitrogen, or oxygen, provided R^{12} is not hydrogen;

a is from 0 and 3;

b is from 0 and 5; and

M is a Group IV to Group VIII metal. The synthesis of the chiral ligands having the formulas IX and X is disclosed in U.S. Provisional Application Nos.

60/236,564 and 60/264,411, both of which are incorporated by reference in their entireties.

The alkyl groups that may be represented by each of R^7 , R^8 , R^9 , R^{10} , R^{11} , and R^{12} in formulas IX and X may be straight- or branched-chain, aliphatic hydrocarbon radicals containing up to 20 carbon atoms and may be substituted, for example, with one to three groups selected from C_1 - C_6 -alkoxy, cyano, C_2 - C_6 -alkoxycarbonyl, C_2 - C_6 alkanoyloxy, hydroxy, aryl and halogen. The terms " C_1 - C_6 -alkoxy," " C_2 - C_6 -alkoxycarbonyl," and " C_2 - C_6 -alkanoyloxy" are used to denote radicals corresponding to the structures $-OR^{13}$, $-CO_2R^{13}$, and $-OCOR^{13}$, respectively, wherein R^{13} is C_1 - C_6 -alkyl or substituted C_1 - C_6 -alkyl. The term " C_3 - C_8 -cycloalkyl" is used to denote a saturated, carbocyclic hydrocarbon radical having three to eight carbon atoms.

The aryl groups for each of R^7 , R^8 , R^9 , R^{10} , R^{11} , and R^{12} in formulas IX and X may include phenyl, naphthyl, or anthracenyl and phenyl, naphthyl, or anthracenyl substituted with one to three substituents selected from C_1 - C_6 -alkyl, substituted C_1 - C_6 -alkyl, C_6 - C_{10} aryl, substituted C_6 - C_{10} aryl, C_1 - C_6 -alkoxy, 5 halogen, carboxy, cyano, C_1 - C_6 -alkanoyloxy, C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulfonyl, trifluoromethyl, hydroxy, C_2 - C_6 -alkoxycarbonyl, C_2 - C_6 -alkanoylamino and $-O-R^{14}$, $S-R^{14}$, $-SO_2-R^{14}$, $-NHSO_2R^{14}$ and $-NHCO_2R^{14}$, wherein R^{14} is phenyl, naphthyl, or phenyl or naphthyl substituted with one to three groups selected from C_1 - C_6 -alkyl, C_6 - C_{10} aryl, C_1 - C_6 -alkoxy and halogen.

10

The heteroaryl radicals include a 5- or 6- membered aromatic ring containing one to three heteroatoms selected from oxygen, sulfur and nitrogen. Examples of such heteroaryl groups are thienyl, furyl, pyrrolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, triazolyl, thiadiazolyl, 15 oxadiazolyl, tetrazolyl, pyridyl, pyrimidyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, indolyl and the like. The heteroaryl radicals may be substituted, for example, with up to three groups such as C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, substituted C_1 - C_6 -alkyl, halogen, C_1 - C_6 -alkylthio, aryl, arylthio, aryloxy, C_2 - C_6 -alkoxycarbonyl and C_2 - C_6 -alkanoylamino. The heteroaryl radicals also 20 may be substituted with a fused ring system, e.g., a benzo or naphtho residue, which may be unsubstituted or substituted, for example, with up to three of the groups set forth in the preceding sentence. The term "halogen" is used to include fluorine, chlorine, bromine, and iodine.

25 In one aspect of the invention, when the chiral ligand is a compound having the formula IX or X, R^{12} is C_1 to C_6 alkyl (e.g., methyl); R^7 is hydrogen or C_1 to C_6 alkyl (e.g., methyl); R^8 is aryl (e.g., phenyl), ethyl, isopropyl, or cyclohexyl; R^9 is aryl (e.g., phenyl); R^{10} and R^{11} are hydrogen; and M is iron, ruthenium, or osmium.

30

The chiral ligand/metal complex can be prepared and isolated, or, in the alternative, it can be prepared *in situ*. The preparation of chiral ligand/metal

complexes is generally known in the art. The chiral ligand to metal molar ratio can be from 0.5:1 to 5:1, preferably from 1:1 to 1.5:1. The amount of chiral ligand/metal complex may vary between 0.0005 and 0.5 equivalents based on compound VI, with more catalyst leading to a faster reaction.

5


The hydrogenation reaction is conducted under an atmosphere of hydrogen, but other materials that are inert to the reaction conditions may also be present. The reaction can be run at atmospheric pressure or at elevated pressure of from 0.5 to 200 atmospheres. The reaction temperature can be varied to modify the rate of conversion, usually between ambient temperature and the boiling point (or apparent boiling point at elevated pressure) of the lowest boiling component of the reaction mixture. In one aspect of the invention, the hydrogenation step is conducted at from -20 °C to 100 °C. The reaction is usually run in the presence of a solvent. Examples of useful solvents include, but are not limited to, aliphatic hydrocarbons such as hexane, heptane, octane and the like, aromatic hydrocarbons such as toluene, xylenes, and the like, cyclic or acyclic ethers such as tert-butyl methyl ether, diisopropyl ether, tetrahydrofuran and the like, dialkyl ketones such as acetone, diethyl ketone, methyl ethyl ketone, methyl propyl ketone and the like, or polar aprotic solvents such as dimethylformamide, dimethyl sulfoxide, N-methylpyrrolidone and the like.

10

15

20

Compounds having the formula VII can be converted to the amide compounds VIII



wherein R¹ and R² are, independently, hydrogen, substituted or unsubstituted, branched or straight chain C₁ to C₂₀ alkyl; substituted or unsubstituted C₃ to C₈ cycloalkyl; substituted or unsubstituted C₆ to C₂₀ aryl; or substituted or unsubstituted C₄ to C₂₀ heteroaryl,

5 n is from 0 to 5, and

the stereochemistry at carbon a is substantially R or S,

10 comprising reacting a compound having the formula VII with NH₄OH.

The amount of NH₄OH can vary, wherein from 1 to 10 equivalents of NH₄OH per 1.0 equivalent of the compound having the formula VII can be used. The reaction is generally performed in water optionally in the presence of
15 a water-miscible organic solvent, including, but not limited to, a lower alcohol such as methanol or ethanol, THF, DMF, or DMSO. The reaction is preferably performed in water as the sole solvent. The reaction temperature can also vary, however; the reaction is typically performed from 0 °C to 50 °C. The compounds having the formula VIII can be used to treat a number of different
20 maladies, some of which are disclosed in U.S. Patent No. 4,696,943, which is incorporated by reference in its entirety.

In summary, the invention provides lactam-substituted propanoic acid derivatives that are useful precursors to enantiomerically-pure lactam-
25 substituted propanoic acid derivatives. The invention provides an efficient method for making the lactam-substituted propanoic acid derivatives as well as the enantiomerically enriched compounds, which will ultimately be used to produce pharmaceutical compounds.

30 Throughout this application, where publications are referenced, the disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains.

EXAMPLES

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how the compositions of matter and methods claimed herein are made and evaluated, and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers (e.g., amounts, temperature, etc.) but some errors and deviations should be accounted for. Unless indicated otherwise, parts are by weight, temperature is in °C or is at room temperature and pressure is at or near atmospheric.

Example 1: Preparation of Methyl 2-(2-Pyrrolidino)propanoate

Step 1: Preparation of Hydroxy-acid 2a (formula II, n=2, R¹=R³=R⁴=H):

Glyoxylic acid monohydrate (20.2 g; 220 mmol; 1.1 equiv) was slurried in 200 mL of diethyl ether. Pyrrolidinone (15.2 mL; 200 mmol) was added and the resulting mixture was stirred at ambient temperature for two days to afford a white precipitate. The precipitate was isolated by filtration, washed with ether, and air-dried to afford 36.06 g (99%) of hydroxy-acid 2a.

¹H NMR (DMSO-d₆) δ 5.55 (s, 1 H); 3.5-3.3 (m, 2H); 3.36 (s, 3H); 2.46 (m, 2H); 2.07 (m, 2H).

Step 2: Preparation of Ether-ester 3a (formula III, n = 2, R¹=H, R³ = R⁴ = Me):

Hydroxy-acid 2a (25.3 g; 160 mmol) was dissolved in methanol (370 mL), cooled in ice-water, and treated with concentrated sulfuric acid (5.0 mL; 0.093 mmol; 0.57 molar equiv). The reaction mixture was allowed to warm to ambient temperature and allowed to stir for 5 days. Solid sodium bicarbonate (17.1g; 203 mmol; 1.27 equiv) was added and the reaction mixture was stirred for 30 min. The reaction mixture was filtered and the filtrate was concentrated. The resulting material was diluted with water (25 mL) and extracted with ethyl acetate (2 x 50 mL). The combined extracts were dried with magnesium sulfate and the solvent was evaporated to afford 22.84 g (76%) of 3a.

^1H NMR (CDCl_3) δ 5.64 (s, 1 H); 3.78 (s, 3H); 3.40 (m, 1H); 3.16 (m, 1H); 2.24 (m, 2H); 1.90 (m, 2H).

Steps 3 and 4: Preparation of Phosphonate Ester 5a (formula V, $n = 2$, $\text{R}^1 = \text{H}$, $\text{R}^3 = \text{R}^6 = \text{Me}$):

Ether-ester 3a (9.36 g; 50 mmol) was dissolved in 50 mL of toluene. The reaction mixture was heated to 70°C and phosphorus trichloride (4.36 mL; 50 mmol; 1.0 equiv) was added. The reaction mixture was heated at 70°C for 18 h to afford chloride 4a (formula IV, $\text{X} = \text{Cl}$, $\text{R}^3 = \text{Me}$). Trimethyl phosphite (5.9 mL; 50 mmol; 1.0 equiv) was then added. The resulting mixture was heated for 36 h and then cooled to ambient temperature. The volatiles were stripped at reduced pressure and the remaining material was dissolved in 100 mL of ethyl acetate. The mixture was washed with saturated aqueous sodium bicarbonate (2 x 25 mL), and the resulting aqueous solution was back-extracted three times with ethyl acetate. The combined ethyl acetate solution was dried with magnesium sulfate and concentrated to afford 4.35 g (33%) of phosphonate 5a.

^1H NMR (CDCl_3) δ 5.46 (d, 1H, $J_{\text{P-H}} = 24.72$ Hz); 3.84 (d, 3H, $J_{\text{P-H}} = 10.99$ Hz); 3.80 (s, 3H); 3.78 (d, 3H, $J_{\text{P-H}} = 11.26$ Hz) 3.8 (m, 1H); 3.66 (m, 1H); 2.42 (m, 2H); 2.08 (m, 2 H).

Step 5: Preparation of Enamide 6a (formula VI, $n = 2$, $\text{R}^1 = \text{H}$, $\text{R}^2 = \text{R}^3 = \text{Me}$):

Phosphonate 5a (2.65 g; 10 mmol) was dissolved in 10 mL of tetrahydrofuran. The mixture was cooled to -78°C and tetramethylguanidine (2.88 mL; 15 mmol; 1.5 equiv) was added. The reaction mixture was stirred for 15 min and acetaldehyde (0.84 mL; 15 mmol; 1.5 equiv) was added. The resulting mixture was stirred at -78°C for 1 h and then warmed to ambient temperature and stirred overnight. The volatiles were evaporated at reduced pressure and water (10 mL) was added. The mixture was extracted three times with ethyl acetate and the combined extracts were dried with magnesium sulfate and concentrated. The crude product was filtered through a pad of flash silica gel and eluted with ethyl acetate to afford 1.37 g (75%) of 6a as a mixture of *E*

and *Z* isomers.

E-6a: ^1H NMR (CDCl_3) δ 7.05 (q, 1H, $J = 7.14$ Hz); 3.74 (s, 3H); 3.55 (t, 2H, $J = 6.86$ Hz); 2.46 (m, 2H); 2.15 (m, 2H); 1.77 (d, 3H, $J = 7.19$ Hz).

Z-6a: ^1H NMR (CDCl_3) δ 5.99 (q, 1H, $J = 7.42$ Hz); 3.78 (s, 3H); 2.02 (d, 3H, $J = 7.42$ Hz).

Step 6: Hydrogenation of Enamide 6a to afford ester 7a (formula VII, $n = 2$, $R^1 = \text{H}$, $R^2 = R^3 = \text{Me}$) using Ligand 9a (formula IX, $R^{12} = R^7 = \text{Me}$, $R^8 = R^9 = \text{Ph}$, $a = b = 0$, $M = \text{Fe}$):

10 A Fischer-Porter tube was charged with ligand 9a (22 mg, 0.036 mmol; 0.064 equiv) and anhydrous THF (4.0 mL) under an argon atmosphere. Argon was bubbled through the solution for 20 minutes before the addition of bis(1,5-cyclooctadienyl)rhodium trifluoromethanesulfonate (12 mg; 0.026 mmol; 0.047 equiv). The solution was stirred at 25°C for 5 minutes or until all bis(1,5-cyclooctadienyl)rhodium trifluoromethanesulfonate had dissolved. Enamide 5a
15 (100 mg, 0.55 mmol) was added *via* syringe. The vessel was capped and pressurized to 40 psi H_2 . After 18 hours, the mixture was diluted with hexane (4.0 mL) and filtered through silica gel to remove the catalyst. The product 7a was isolated as an oil (99% yield, 96.2% ee as determined by chiral GC).
20 ^1H NMR (CDCl_3) δ 4.71-4.65 (m, 1H), 3.71 (s, 3H), 3.55-3.47 (m, 1H), 3.38-3.30 (m, 1H), 2.45-2.40 (t, $J=8.4$ Hz, 2H), 2.12-1.97 (m, 3H), 1.74-1.64 (m, 1H), 0.94-0.89 (t, $J=7.5$ Hz, 3H). Chiral GC analysis [Cyclosil-B (J&W Scientific), 40°C for 4 min to 175°C at 70°C/min, hold at 175°C for 12 minutes: $t_R = 23.19$ (major enantiomer), $t_R = 23.24$ (minor enantiomer)].

25

Example 2:

In Situ Preparation of Compound VI ($n=2$, $R^1 = \text{H}$, $R^2 = R^3 = \text{Me}$):

Ether-ester 3a (9.36 g; 50 mmol) was dissolved in 25 mL of toluene.

30 The reaction mixture was heated to 50°C and phosphorus trichloride (4.4 mL; 50 mmol; 1.0 equiv) was added and the mixture was held at 50°C for 16 h. The

reaction mixture was cooled to ambient temperature and sodium bicarbonate (10.5 g; 125 mmol; 2.5 equiv) was added. The mixture was stirred for 15 min, filtered, and the volatiles were stripped. The solution was reconstituted by the addition of toluene (25 mL) and heated to 70°C. Trimethyl phosphite (6.5 mL; 55 mmol; 1.1 equiv) was added and the reaction mixture was heated at 70°C for 24 h to completely consume 4a. The mixture was then cooled to 2°C and acetaldehyde (4.2 mL; 75 mmol; 1.5 equiv) was added. Tetramethylguanidine (9.41 mL; 75 mmol; 1.5 equiv) was then added slowly dropwise with an attendant exotherm to 8.5°C. The reaction mixture was allowed to warm to ambient temperature and stirred overnight to consume phosphonate 5a. Water (15 mL) was added and the layers were separated. The aqueous layer was extracted with ethyl acetate (2 x 15 mL) and the organic solutions were dried and concentrated to afford a total of 9.23 g of crude 6a. This material was filtered through a pad of flash silica gel and eluted with ethyl acetate to afford 7.13 g (78% overall from 3a) of 6a.

Example 3: Preparation of Ethyl 2-(2-Pyrrolidino)propanoate

Step 2: Preparation of Ether-ester 3b (formula III, n = 2, R¹=H, R³ = R⁴ = Et):

Hydroxy-acid 2a (7.96 g; 50 mmol) was dissolved in ethanol (25 mL), and triethyl orthoformate (18.3 mL; 110 mmol; 2.2 equiv) was added. p-Toluenesulfonic acid (0.48 g; 2.5 mmol; 0.05 equiv) was added and the reaction mixture was heated to 60°C for 24 h. Solid sodium bicarbonate (0.50 g; 6 mmol; 0.12 equiv) was added and the reaction mixture was stirred for 15 min. The volatiles were evaporated at reduced pressure and the remaining material was dissolved in 1:1 toluene:ethyl acetate (25 mL), filtered, and concentrated to afford 10.25 g (95%) of 3b.

¹H NMR (CDCl₃) δ 5.72 (s, 1 H); 4.24 (q, 2H, J = 7.14 Hz); 3.57 (q, 2H, J = 6.87 Hz); 3.5-3.3 (m, 2H); 2.47 (t, 2H, J = 7.42 Hz); 2.06 (m(5), 2H, J = 7.42 Hz); 1.29 (t, 3H, J = 7.14 Hz); 1.25 (t, 3H, J = 6.87 Hz).

Steps 3 and 4: Preparation of Phosphonate Ester 5b (formula V, $n = 2$, $R^1 = H$, $R^3 = Et$, $R^6 = Me$):

Ether-ester **3b** (5.38 g; 25 mmol) was dissolved in 12.5 mL of toluene. The reaction mixture was heated to 50°C and phosphorus trichloride (2.2 mL; 25 mmol; 1.0 equiv) was added. The reaction mixture was heated at 50°C for 12 h and then at 70°C for 24 h to completely consume **3b** and afford chloride **4b** (formula IV, $X = Cl$, $R^3 = Et$). Solid sodium bicarbonate (5.25 g; 62.5 mmol; 2.5 equiv) was added and the mixture was stirred for 15 min. The reaction mixture was then filtered and concentrated at reduced pressure. Toluene (12.5 mL) was added and the reaction mixture was heated to 70°C where trimethyl phosphite (3.24 mL; 25 mmol; 1.0 equiv) was added. The resulting mixture was heated at 70°C for 24 h and then cooled to ambient temperature. The volatiles were evaporated at reduced pressure to afford 7.00 g (99%) of phosphonate **5b**, which was used without further purification.

1H NMR ($CDCl_3$) δ 5.43 (d, 1H, $J_{P-H} = 24.72$ Hz); 4.25 (q, 2H, $J = 7.14$ Hz); 3.84 (d, 3H, $J_{P-H} = 10.99$ Hz); 3.77 (d, 3H, $J_{P-H} = 10.99$ Hz); 3.8 (m, 1H); 3.66 (m, 1H); 2.40 (m, 2H); 2.06 (m, 2H); 1.29 (t, 3H, $J = 7.14$ Hz).

Step 5: Preparation of Enamide 6b (formula VI, $n = 2$, $R^1 = H$, $R^3 = Et$, $R^2 = Me$):

Phosphonate **5b** (7.00 g; 25 mmol) was dissolved in 12.5 mL of toluene. The mixture was cooled to 2°C and acetaldehyde (2.1 mL; 37.5 mmol; 1.5 equiv) was added. Tetramethylguanidine (3.5 mL; 27.5 mmol; 1.1 equiv) was added dropwise accompanied by a moderate exotherm. The reaction mixture was allowed to warm to ambient temperature and stirred overnight and then heated to 50°C for 12 h to completely consume **5b**. The reaction mixture was diluted with 3 N HCl (10 mL) and the layers were separated. The aqueous layer was extracted twice with ethyl acetate and the combined organic solution was dried with magnesium sulfate and concentrated. The crude product was filtered through a pad of flash silica gel and eluted with ethyl acetate to afford 2.85 g (58%) of **6b** as a mixture of *E* and *Z* isomers.

E-6b: 1H NMR ($CDCl_3$) δ 7.05 (q, 1H, $J = 7.14$ Hz); 4.20 (q, 2H, $J = 7.14$ Hz); 3.55 (t, 2H, $J = 7.14$ Hz); 2.48 (t, 2H, $J = 7.69$ Hz); 2.15 (m(5), 2H, $J = 7.69$

Hz); 1.77 (d, 3H, J = 7.14 Hz); 1.28 (t, 3H, J = 7.14 Hz).

Z-6b: ^1H NMR (CDCl_3) δ 6.0 (q, 1H); 3.78 (s, 3H); 4.12 (q, 2H); 1.95 (d, 3H).

Step 6: Hydrogenation of Enamide 6b to afford ester 7b (formula VII, n = 1, R¹=H, R³ = Et, R² = Me) using Ligand 10a (formula X, R¹² = R⁷ = Me, R⁸ = R⁹ = Ph, a = b = 0, M = Fe): Bis(1,5-cyclooctadienyl)rhodium trifluoromethanesulfonate (5 μmol , 2.3 mg) was placed into a reaction vessel and purged with argon for 20 min. A solution of 10a (6 μmol , 3.7 mg) in anhydrous THF (2.0 mL) was degassed with Ar for 20 minutes, then added via cannula to the bis(1,5-cyclooctadienyl)rhodium trifluoromethanesulfonate. This solution was stirred at 25°C under Ar for 20 minutes. A solution of enamide 6b (0.5 mmol) in anhydrous THF (2.0 mL) was degassed with Ar for 20 minutes, then added to the catalyst solution via cannula. The solution was then flushed with H₂ and pressurized to 20 psig H₂. A sample was taken after 1 hour and analyzed for enantiomeric excess using standard chiral gas chromatography. 7b was formed in 95.0% ee with >99% conversion. Chiral GC analysis [Cyclosil-B (J&W Scientific), 30 m x 0.25 mm ID, film thickness 0.25 μm , 60-140°C 40°C/min, 140°C 30 min, 20 psig He: t_R(minor enantiomer) 26.26 min, t_R(major enantiomer) 26.82 min.

20

Example 4: Preparation of Amide

Step 1: Conversion of Ester 7a (formula VII, n = 1, R¹=H, R² = R³ = Me) to Amide 8 (formula VIII, n = 1, R¹=H, R² = Me):

Ester 7a (93.6% ee; 0.93 g; 5.0 mmol) was combined with ammonium hydroxide (28% NH₃; 1.0 mL; 15 mmol; 3 equiv) and stirred at ambient temperature overnight, at which time tlc analysis indicated no 7a. The mixture was diluted with water (5 mL) and extracted with three 5 mL portions of dichloromethane. The combined extracts were dried with sodium sulfate and concentrated to afford 0.62 g (73%) of 8 which possessed 92.6% ee according to chiral GC analysis, indicating very little loss of enantiomeric purity. The crude

30

product was recrystallized from hot acetone (3.1 mL; 5 mL/g) by cooling to 0°C.

The resulting crystals were collected by filtration, washed with 1:1 acetone:heptane, and air-dried to afford 0.37 g (43%) of **8** which was 99.6% ee according to chiral GC.

- 5 ¹H NMR (CDCl₃) δ 6.43 (br s, 1H); 5.72 (br s, 1 H); 4.455 (dd, 1H, J = 9.07, 6.87 Hz); 3.38 (m, 2H); 2.40 (m, 2H); 2.1-1.9 (m, 3H); 1.67 (m, 1H); 0.886 (t, 3H, J = 7.42 Hz). Chiral GC analysis [Cyclosil-B (J&W Scientific), 175°C for 25 minutes, 15 psi He]: t_R = 21.31 (*S* enantiomer), t_R = 22.06 (*R* enantiomer)]. [α]_D²² -86.7° (c 0.98, acetone).

10


Comparative Example: Conversion of Ester **7a** (formula VII, n = 1, R¹=H, R²=R³ = Me) to Amide **8** (formula VIII, n = 1, R¹=H, R² = Me):

Ester **7a** (93.6% ee; 0.93 g; 5.0 mmol) was dissolved in 2 mL of methanol and treated with ammonium hydroxide (28% ammonia; 1.0 mL; 15 mmol; 3 equiv).

- 15 The mixture was stirred at ambient temperature overnight, at which time tlc analysis indicated some residual **7a**. Additional ammonium hydroxide (1.0 mL; 15 mmol; 3 equiv) was added and the reaction mixture was stirred for 2 days to completely consume **7a**. The reaction mixture was evaporated under reduced pressure and azeotroped several times with toluene under reduced pressure to
20 afford 0.95 g of **8** as a brown oil which was 89.4% ee according to chiral GC analysis, indicating substantial loss of enantiomeric purity.

What is claimed is:

1. A compound having the formula VII



wherein R^1 , R^2 , and R^3 are, independently, hydrogen, a substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl group; a substituted or unsubstituted C_3 to C_8 cycloalkyl group; a substituted or unsubstituted C_6 to C_{20} aryl group; or substituted or unsubstituted C_4 to C_{20} heteroaryl group having at least one heteroatom,


n is from 0 to 5, and

the stereochemistry at carbon a is substantially R or S.

2. The compound of Claim 1, wherein the stereochemistry at carbon a is substantially S.
3. The compound of Claim 1, wherein the heteroatom of the heteroaryl group is oxygen, sulfur, or nitrogen, and the substituent on the substituted alkyl, aryl, or heteroaryl group comprises alkyl, aryl, hydroxy, alkoxy, fluoro, chloro, bromo, iodo, nitro, cyano, or an ester.
4. The compound of Claim 1, wherein R^2 and R^3 are independently selected from methyl or ethyl.
5. The compound of Claim 2, wherein R^1 is hydrogen, R^2 is methyl, R^3 is ..

methyl or ethyl, and n is 2.

6. A method for producing the compound of Claim 1, comprising hydrogenating an enamide having the formula VI




wherein R^1 , R^2 , R^3 , and n are as set forth in Claim 1,

with hydrogen in the presence of a catalyst comprised of a chiral ligand/metal complex to asymmetrically hydrogenate the carbon-carbon double bond of the enamide.

7. The method of Claim 6, wherein the chiral ligand of the chiral ligand/metal complex comprises a phosphine or a bis-phosphine compound and the metal of the chiral ligand/metal complex comprises rhodium, ruthenium, or iridium.
8. The method of Claim 6, wherein the chiral ligand of the chiral ligand/metal complex comprises a phosphine or a bis-phosphine compound and the metal of the chiral ligand/metal complex comprises rhodium.
9. The method of Claim 6, wherein the chiral ligand of the chiral ligand/metal complex comprises a substantially enantiomerically pure bis-phosphine compound comprising a substantially enantiomerically pure chiral backbone linking two phosphine residues, wherein one of the phosphine residues has three phosphorus-carbon bonds and the other phosphine residue has two

phosphorus-carbon bonds and one phosphorus-nitrogen bond wherein the nitrogen is part of the chiral backbone.

10. The method of Claim 6, wherein the chiral ligand of the chiral ligand/metal complex comprises a compound having the formula IX or X



where R^7 , R^8 , R^9 , R^{10} , R^{11} , and R^{12} are, independently, hydrogen, substituted or unsubstituted branched or straight chain C_1 to C_{20} alkyl, substituted or unsubstituted C_3 to C_8 cycloalkyl, substituted or unsubstituted C_6 to C_{20} aryl, and substituted or unsubstituted C_4 to C_{20} heteroaryl, where the heteroatoms are chosen from sulfur, nitrogen, or oxygen, provided R^{12} is not hydrogen;

a is from 0 and 3;


b is from 0 and 5; and

M is a Group IV to Group VIII metal.

11. The method of Claim 10, wherein M comprises iron, ruthenium, or osmium.
12. The method of Claim 10, wherein a and b are 0, R^7 and R^{12} are methyl, R^8 and R^9 are phenyl, and M is iron.
13. The method of Claim 6, wherein the chiral ligand of the chiral ligand/metal complex comprises the substantially pure enantiomer or diastereomer of 2,3-*O*-isopropylidene-2,3-dihydroxy-1,4-bis(diphenylphosphino)butane; 2,2'-

bis(diphenylphosphino)-1,1'-binaphthyl; 1,2-*bis*-2,5-dialkylphospholano(benzene); 1,2-*bis*-2,5-dialkylphospholano(ethane); 2,3-*bis*-(diphenylphosphino)butane; or 2-diphenylphosphinomethyl-4-diphenylphosphino-1-*t*-butoxycarbonylpyrrolidine.

14. The method of Claim 6, wherein the metal of the chiral ligand/metal complex is from 0.0005 to 0.5 equivalents per 1.0 equivalent of the compound having the formula VI.
15. The method of Claim 6, wherein the hydrogenation step is conducted under an atmosphere of hydrogen at from 0.5 to 200 atmospheres.
16. The method of Claim 6, wherein the hydrogenation step is conducted in a solvent comprising an aliphatic hydrocarbon, an aromatic hydrocarbon, a cyclic ether, an acyclic ether, a halogenated hydrocarbon, a dialkyl ketone, a polar aprotic solvent, or a combination thereof.
17. The method of Claim 6, wherein the hydrogenation step is conducted at from -20 °C to 100 °C.
18. A compound having the formula XI




wherein R^1 , R^3 , and R^4 are, independently, hydrogen, a substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl group; a substituted or unsubstituted C_3 to C_8 cycloalkyl group; a substituted or unsubstituted C_6

to C₂₀ aryl group; or substituted or unsubstituted C₄ to C₂₀ heteroaryl group,
and

n is from 0 to 5.


19. The compound of Claim 18, wherein n is 2.
20. The compound of Claim 18, wherein R¹, R³, and R⁴ are hydrogen and n is 2.
21. The compound of Claim 18, wherein R¹ is hydrogen, R³ and R⁴ are methyl,
and n is 2.
22. A method for producing a compound having the formula II,



wherein R¹ is hydrogen, substituted or unsubstituted, branched or straight
chain C₁ to C₂₀ alkyl; substituted or unsubstituted C₃ to C₈ cycloalkyl;
substituted or unsubstituted C₆ to C₂₀ aryl; or substituted or unsubstituted C₄
to C₂₀ heteroaryl, and


n is from 0 to 5,

comprising reacting a compound having the formula I



with glyoxylic acid.


23. The method of Claim 22, wherein the glyoxylic acid is present in the amount from 0.8 to 2 equivalents per 1.0 equivalent of the compound having the formula I.
24. A method for producing the compound having the formula III,



wherein R¹, R³, and R⁴ are, independently, a substituted or unsubstituted, branched or straight chain C₁ to C₂₀ alkyl group; a substituted or unsubstituted C₃ to C₈ cycloalkyl group; a substituted or unsubstituted C₆ to C₂₀ aryl group; or a substituted or unsubstituted C₄ to C₂₀ heteroaryl group, wherein R¹ can also be hydrogen, and


n is from 0 to 5,

comprising reacting the compound having formula II



with an alcohol comprising an alkyl alcohol, an aryl alcohol, or a heteroaryl alcohol, wherein the alkyl alcohol is substituted or unsubstituted, branched or straight chain C₁ to C₂₀ alkyl or substituted or unsubstituted C₃ to C₈ cycloalkyl; the aryl alcohol is substituted or unsubstituted C₆ to C₂₀ aryl; and the heteroaryl alcohol is substituted or unsubstituted C₄ to C₂₀ heteroaryl, wherein the heteroatom is oxygen, nitrogen, or sulfur.

25. The method of Claim 24, wherein the alcohol is a C₁ to C₅ alcohol.
26. The method of Claim 24, wherein the alcohol is methanol or ethanol.
27. The method of Claim 24, wherein the alcohol is present in the amount from 2.0 to 5.0 equivalents per 1.0 equivalent of the compound having the formula II.
28. The method of Claim 24, further comprising a dehydrating agent.
29. A compound having the formula V




wherein R^1 and R^3 are, independently, substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl; substituted or unsubstituted C_3 to C_8 cycloalkyl; substituted or unsubstituted C_6 to C_{20} aryl; or substituted or unsubstituted C_4 to C_{20} heteroaryl, wherein R^1 can also be hydrogen,

R^6 is substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl or substituted or unsubstituted C_3 to C_8 cycloalkyl, and

n is from 0 to 5.

30. The compound of Claim 29, wherein n is 2 and R^1 is hydrogen.
31. The compound of Claim 30, wherein R^3 is methyl or ethyl.
32. The compound of Claim 31, wherein R^6 is methyl or ethyl.
33. A method of producing the compound of Claim 29, comprising reacting a compound having the formula IV




wherein R^1 and R^3 are, independently, substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl; substituted or unsubstituted C_3 to C_8 cycloalkyl; substituted or unsubstituted C_6 to C_{20} aryl; or substituted or unsubstituted C_4 to C_{20} heteroaryl, wherein R^1 can also be hydrogen,

X is fluoride, chloride, bromide, or iodide, and

n is from 0 to 5,

with a phosphite having the formula $P(OR^6)_3$, wherein R^6 is substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl or substituted or unsubstituted C_3 to C_8 cycloalkyl.

34. The method of Claim 33, wherein X is chloride or bromide.
35. The method of Claim 33, wherein R^6 is methyl or ethyl.
36. The method of Claim 33, wherein the phosphite is present in the amount from 0.8 to 1.2 equivalents per 1.0 equivalent of the compound having the formula IV.
37. A method of producing a compound having the formula IV,




wherein R^1 and R^3 are, independently, substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl; substituted or unsubstituted C_3 to C_8 cycloalkyl; substituted or unsubstituted C_6 to C_{20} aryl; or substituted or unsubstituted C_4 to C_{20} heteroaryl, wherein R^1 can also be hydrogen,

X is fluoride, chloride, bromide, or iodide, and

n is from 0 to 5,

comprising reacting a compound having the formula III




wherein R^1 , R^3 , and R^4 are, independently, a substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl group; a substituted or unsubstituted C_3 to C_8 cycloalkyl group; a substituted or unsubstituted C_6 to C_{20} aryl group; or a substituted or unsubstituted C_4 to C_{20} heteroaryl group, wherein R^1 can also be hydrogen, and

n is from 0 to 5,

with a compound having the formula PX_3 , wherein X is fluoro, chloro, bromo, or iodo.

38. A compound having the formula VI




wherein R^1 , R^2 , and R^3 are, independently, hydrogen, substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl; substituted or

unsubstituted C₃ to C₈ cycloalkyl; substituted or unsubstituted C₆ to C₂₀ aryl;
or substituted or unsubstituted C₄ to C₂₀ heteroaryl, and

n is from 0 to 5.

39. The compound of Claim 38, wherein n is 2 and R¹ is hydrogen.
40. The compound of Claim 39, wherein R² and R³ are methyl.
41. The compound of Claim 39, wherein R² is methyl and R³ is ethyl.
42. A method for producing the compound of Claim 38, comprising reacting a compound having the formula V



wherein R¹ and R³ are, independently, substituted or unsubstituted, branched or straight chain C₁ to C₂₀ alkyl; substituted or unsubstituted C₃ to C₈ cycloalkyl; substituted or unsubstituted C₆ to C₂₀ aryl; or substituted or unsubstituted C₄ to C₂₀ heteroaryl, wherein R¹ can be hydrogen,


R⁶ is substituted or unsubstituted, branched or straight chain C₁ to C₂₀ alkyl or substituted or unsubstituted C₃ to C₈ cycloalkyl, and

n is from 0 to 5,

with an aldehyde having the formula HC(O)R^2 , wherein R^2 is hydrogen, substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl; substituted or unsubstituted C_3 to C_8 cycloalkyl; substituted or unsubstituted C_6 to C_{20} aryl; or substituted or unsubstituted C_4 to C_{20} heteroaryl

in the presence of a base.

43. The method of Claim 42, wherein the base comprises an amidine base or a guanidine base.
44. The method of Claim 42, wherein the base comprises 1,5-diazabicyclo(4.3.0)non-5-ene; 1,8-diazabicyclo(5.4.0)undec-7-ene, or tetramethylguanidine.
45. The method of Claim 42, wherein the base is present in the amount from 1.0 to 2.0 equivalents per 1.0 equivalent of the compound having the formula V.
46. The method of Claim 42, wherein the aldehyde is present in the amount from 0.8 to 1.5 equivalents per 1.0 equivalent of the compound having the formula V.
47. The method of Claim 42, wherein the aldehyde is acetaldehyde.
48. A method for producing the compound of Claim 38 *in situ*, comprising
 - (a) reacting a compound having the formula III



wherein R^3 and R^4 are, independently, hydrogen, substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl group; substituted or unsubstituted C_3 to C_8 cycloalkyl group; substituted or unsubstituted C_6 to C_{20} aryl group; or substituted or unsubstituted C_4 to C_{20} heteroaryl group, wherein R^1 can also be hydrogen


with PX_3 , wherein X is fluoride, chloride, bromide, or iodide, to produce a halogenated lactam;

- (b) reacting the halogenated lactam produced in step (a) with a phosphite having the formula $P(OR^6)_3$, wherein R^6 is substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl or substituted or unsubstituted C_3 to C_8 cycloalkyl, to produce a phosphonated lactam; and
- (c) reacting the phosphonated lactam produced in step (b) with an aldehyde having the formula $HC(O)R^2$, wherein R^2 is hydrogen, substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl; substituted or unsubstituted C_3 to C_8 cycloalkyl; substituted or unsubstituted C_6 to C_{20} aryl; or substituted or unsubstituted C_4 to C_{20} heteroaryl,

in the presence of a base,

wherein steps (a), (b), and (c) are performed *in situ*.

49. A method for producing a compound having the formula VIII




wherein R^1 and R^2 are, independently, hydrogen, substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl; substituted or unsubstituted C_3 to C_8 cycloalkyl; substituted or unsubstituted C_6 to C_{20} aryl; or substituted or unsubstituted C_4 to C_{20} heteroaryl,

n is from 0 to 5, and

the stereochemistry at carbon a is substantially R or S,

comprising reacting a compound having the formula VII



wherein R^3 is substituted or unsubstituted, branched or straight chain C_1 to C_{20} alkyl; substituted or unsubstituted C_3 to C_8 cycloalkyl; substituted or unsubstituted C_6 to C_{20} aryl; or substituted or unsubstituted C_4 to C_{20} heteroaryl,

with NH_4OH .

50. The method of Claim 49, wherein the NH_4OH is present in the amount from 1 to 10 equivalents per 1.0 equivalent of the compound having the formula VII.